

The BeppoSAX High Energy Large Area Survey (HEL-LAS): a progress report

A. Comastri¹, F. Fiore^{2,3,4}, C. Vignali^{1,5}, F. La Franca⁶, G. Matt⁶

¹*Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italy*

²*Osservatorio Astronomico di Roma, Via Frascati 33, I-00044 Monteporzio, Italy*

³*BeppoSAX Science Data Center, Via Corcolle 19, I-00131 Roma, Italy*

⁴*Center of Astrophysics, 60 Garden Street, Cambridge MA 02138 USA*

⁵*Dipartimento di Astronomia, via Ranzani 1, I-40127 Bologna, Italy*

⁶*Dipartimento di Fisica, Università degli Studi "Roma Tre", Via della Vasca Navale 84, I-00146 Roma, Italy*

Abstract.

The integrated emission of highly obscured AGN is expected to provide a major contribution to the X-ray energy density in the Universe: the X-ray background (XRB). The study of these objects is possible only at energies where the effects of absorption are less severe. For this reason we have carried out the BeppoSAX High Energy LLarge Area Survey in the hardest band (5–10 keV) accessible so far with imaging X-ray instruments. The source surface density at the survey limiting flux accounts for a significant fraction (20–30 %) of the hard XRB. The X-ray data complemented by multiwavelength follow-up observations suggest that a large fraction of the hard sources are AGN and that X-ray absorption with column densities in the range $10^{22-23.5} \text{ cm}^{-2}$ is common among them. The great diversity in their optical-near-IR properties suggests that the optical appearance of obscured sources is a function of the X-ray luminosity. We briefly discuss the implications of these findings for the XRB models.

1 Introduction

The identification breakdown of soft X-ray surveys has clearly established that AGN are by far the dominant population of the X-ray sky (see [12] for the most recent update) implying that about 60–70 % of the soft XRB has been already resolved into AGN at a limiting 0.5–2 keV flux of $\sim 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$. Thanks to the imaging capabilities of ASCA and BeppoSAX detectors the number of hard X-ray selected optically identified objects is increasing and several dozens of identifications are now available confirming that most of these sources are indeed AGN ([8], [1]). Their contribution to the 2–10 keV XRB is of the order of 20–25 % at a flux limit of $\sim 5 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$.

The present findings provide support to the XRB synthesis models which, in the framework of the unified scheme, assume that a mixture of unabsorbed and

absorbed AGN can account for almost the entire XRB spectral intensity in the 2–100 keV energy range ([17], [4], [10], [15]) Even though the so far proposed models differ in several of the assumptions concerning the AGN luminosity function and the X-ray spectral properties, they all agree in predicting that the fraction of obscured AGN is rapidly increasing towards high energy and faint fluxes. It turns out that these sources can be efficiently discovered with sensitive hard X-ray surveys.

In order to quantitatively test AGN synthesis models for the XRB we have carried out an X-ray survey in the hardest band accessible with the present imaging detectors: the 5–10 keV *BeppoSAX* High Energy LLarge Area Survey (HELLAS).

2 Source Counts

About 80 square degrees of sky have been surveyed in the 5–10 keV band using several *BeppoSAX* MECS ([2]) high Galactic latitude ($|b| > 20$ deg) fields. All MECS pointings cover different sky positions. The fields were selected among public data (as March 1999) and our proprietary data excluding those fields centered on extended sources and bright Galactic objects. A robust detection algorithm has been used on the coadded MECS1 MECS2 and MECS3 (or MECS2 plus MECS3 after the failure of MECS1 in May 1997) images and the quality of the detection has been checked interactively for all the 147 sources of the final sample. Background subtracted count rates were converted to 5–10 keV fluxes assuming a power law spectrum with energy index $\alpha_E = 0.6$. A detailed description of the survey and the detection procedure is presented in [9].

The 5–10 keV logN–logS of the HELLAS sources is reported in Figure 1. The source surface density at the survey limit (4.8×10^{-14} erg cm $^{-2}$ s $^{-1}$) is 17 ± 6 sources per square degree. The error bars on the binned integral counts account for both the statistic and systematic uncertainties, the latter due to the lack of information on the intrinsic spectrum of the faint sources. The systematic error has been estimated assuming a range of spectral shapes ($0.2 < \alpha_E < 1.0$) to convert count rates into fluxes. The HELLAS counts are in very good agreement with the recent *ASCA* results ([3], [18]) obtained in the 2–10 keV band (Figure 1). Given that both the HELLAS and the *ASCA* counts have been computed assuming the same prescription for the spectral slope the comparison is straightforward. The dashed line in Figure 1 corresponds to the best-fit to the *ASCA* counts computed by [7] and converted in the 5–10 keV band with $\alpha_E = 0.6$. The cumulative flux of the HELLAS sources accounts for a significant fraction (20–30 %) of the 5–10 keV XRB. The main uncertainty on the resolved fraction is due to the still poorly understood normalization of the extragalactic XRB spectrum as measured by the different satellites (see [5] and [19] for more details).




Figure 1: The 5–10 keV HELLAS counts compared with the ASCA 2–10 keV counts (dashed line). A power law spectrum with $\alpha_E = 0.6$ has been adopted for the flux conversion.

3 Hardness ratios

The HELLAS sources are too faint to perform a spectral fit. In order to study their spectral properties we have computed for each source, whenever possible, two X-ray colours defined as $HR1 = (M-L)/(M+L)$ and $HR2 = (H-M)/(H+M)$ where H, M and L are the number of counts in the H=4.5–10 keV M=2.5–4.5 keV and L=1.3–2.5 keV energy ranges. A wide range of spectral properties is evident from the analysis of the color-color diagram reported in Figure 2. For example extremely hard sources, with nearly all the photons detected only above 4.5 keV, populate the upper right part of the diagram. The X-ray colours depend from the intrinsic spectrum and from the source redshift. The colours expected for a range of spectral shapes at different redshifts has been computed following the same procedure adopted by [6] for their ASCA survey.

The big star at $HR1 = 0.25$ and $HR2 = 0$ represents an unabsorbed power law spectrum with $\alpha_E = 0.4$. The rightmost solid curve connecting the open squares indicates the colours for an AGN-like power law ($\alpha_E = 0.8$) absorbed by an increasing value of the column density : $\log N_H = 0, 22, 22.7, 23, 23.7, 24$ at $z=0$. The dashed curve is the same but at $z=0.4$. The innermost solid curves have been computed at $z=0$ assuming the same model but allowing a fraction of 10% and 1% respectively to be unabsorbed. These models should be considered as indicative




Figure 2: The X-ray color-color diagram for the 58 sources detected above 3.8σ (see text for details). The optically identified objects are reported with filled symbols.

and indeed they do not cover the entire diagram. As first noted by [6], more complicate spectral shapes, such as those characterizing the sources in the upper left portion of the plot, might be present. ASCA follow-up pointed observations of two relatively bright HELLAS sources [20] allowed to collect enough counts to perform a more detailed spectral analysis. In both of the cases the data are consistent with a hard absorbed power law spectrum making us confident on the robustness of the hardness ratio results.

4 Optical Identifications

The cross-correlation of the HELLAS sample with various source catalogs provided 25 coincidences (19 AGN: 7 radio-loud, 12 radio quiet and 6 clusters of galaxies). In addition, optical spectroscopic follow-up observations have been performed and 22 new identifications (18 AGN) are available ([8], [11]). A detailed discussion of the optical identifications is beyond the purposes of this paper and will be presented by [11].

The average X-ray spectrum as inferred from the softness ratio value $S-H/S+H$ (where S is the number of counts in the 1.3–4.5 keV energy range) is plotted in Figure 3 versus the source redshift and optical classification.

The most important results are the following:

- The fraction of type 2 objects (including in this class Seyfert types 1.8–1.9–2.0 and quasars with a red optical continuum) is of the order of 40–45 %. This percentage is higher than in other optically identified samples of X-ray selected sources from ROSAT and ASCA surveys.
- The degree of obscuration as inferred from the hardness ratio analysis described in the previous section seems to be uncorrelated with the optical reddening indicators, such as the optical line widths and line ratios.




Figure 3: The softness ratio versus redshift for the so far identified sources. Different symbols mark different sources: open circles = broad line quasars with a blue optical continuum, stars = broad lined quasars with a red optical continuum, filled circles = AGN of types 1.8–1.9–2.0, filled squares = emission line galaxies, open triangles = radio-loud AGN open squares = clusters of galaxies. The two dotted lines represent the expected softness ratio for a power law with $\alpha_E=0.4$ and $\alpha_E=0.8$. Dashed lines show the softness ratios of absorbed power law models (for $\alpha_E = 0.8$ and $\log N_H = 22.7, 23, 23.7$, from top to bottom) with the absorber at the source redshift.

- The softness ratio of a few high luminosity broad lined quasars with blue optical colours implies X-ray absorption by a substantial column density (log

$N_H > 23$).

- Optical and near-infrared photometry of ten HELLAS sources carried out at the Italian National Telescope Galileo (TNG) indicates that the optical colors of type 1.8–2.0 and red AGN are dominated by the host galaxy, though the obscured AGN contributes to some of the infrared emission ([14], [21], [20]).

5 A quick comparison with XRB synthesis models

The observed 5–10 keV $\log N$ – $\log S$ is compared with the AGN number counts predictions [4] in the same energy range (Figure 4). Given that in the HELLAS band a column as high as $\log N_H = 23$ is needed to significantly reduce the photon flux, the contribution of sources with a different degree of obscuration is reported splitted into 3 classes : relatively unobscured $\log N_H < 23$, obscured Compton thin $23 < \log N_H < 24$ and almost Compton thick $\log N_H > 24$. According to the model predictions relatively unobscured sources outnumber absorbed objects. The expected fraction of Compton thin sources ranges from 25 to 35 % while the number of Compton thick sources is always negligible. Given that about one third of the AGN reported in Figure 3 have a softness ratio value consistent with a column density $23 < \log N_H < 24$ and none is Compton thick the agreement with the model predictions is remarkable. These findings are at variance with the relatively high space density of Compton thick AGN in the local Universe [13], [16]. We note, however, that the fraction of Compton thick objects has been estimated only for optically selected, nearby Seyferts and thus might not be representative of the more distant X-ray selected population.

6 Discussion

The hard X-ray sky, as surveyed by BeppoSAX, is dominated by AGN with a wide range of X-ray and optical properties. At first glance the present data are in line with what expected from the standard synthesis models for the XRB [4]. In addition, new trends characterizing the AGN population are emerging.

The large dispersion in the hardness/softness ratios suggests that the spectral properties of the HELLAS sources cannot be explained only with a distribution of absorbing column densities as assumed in the XRB models. In agreement with ASCA findings [6], the presence of additional soft X-ray emission above the hard absorbed component is required. The “soft excess” is likely to be due either to a fraction of the nuclear component scattered in the line of sight or to an incomplete covering of the central radiation or to thermal emission originating in a starburst region and/or due to X-ray binaries and supernova remnants. The “soft excess” intensity, which ranges from a few percent to 10–20 % of the total X-ray flux, is not relevant as far as the fit to the hard XRB is concerned, however it should be consistently taken into account when model predictions are compared to hard and




Figure 4: The 5–10 keV counts compared with the prediction of XRB synthesis models ([5]). The solid line represents the total AGN contribution, the other lines the number counts of AGN with various degrees of intrinsic obscuration: $\log N_H < 23$ (dotted), $23 < \log N_H < 24$ (dashed), $\log N_H > 24$ (dot-dashed)

soft X-ray counts. Indeed, if soft excesses are common among obscured AGN, as the HELLAS results indicate, these sources might well be detectable in the soft X-ray band even in the presence of substantial obscuration simply because the flux limit of the ROSAT soft X-ray surveys is about a factor 30–40 deeper than the limit actually reached in the hard X-ray band.

Another interesting and unexpected result concerns the optical and near-infrared properties of the hard X-ray selected sources. Even though detailed studies have been carried out only for a small HELLAS subsample the results do clearly indicate that:

- The broad band optical (U,B,V,R,I) and near-infrared (J, K) colors of HELLAS AGN, spectroscopically classified as type 1.8–2.0 Seyferts or red quasars, are undistinguishable from those of normal passive galaxies.
- There is an increasing evidence that the X-ray absorption properties and the optical appearance of AGN change with redshift and/or luminosity, suggesting that high luminosity, highly obscured quasars are present among optically broad lined blue objects.

If confirmed by future observations, these findings would imply that the X-ray obscured AGN responsible for a large fraction of the hard XRB energy density could be “hidden” among objects which would be optically classified either as normal galaxies or as “normal” blue quasars. It is also worth noting that the

redshift of obscured AGN could be in principle obtained by deep photometric observations of their host galaxies.

A step forward in the study of the sources of the hard XRB will be achieved by the foreseen **XMM** and **Chandra** surveys. In particular the X-ray data will allow to obtain an unbiased estimate of the absorption distribution (a key parameter of the XRB models), while optical and near-infrared follow-up observations are likely to provide new insights on the nature of the absorbing gas (such as the dust-to-gas ratio) and on the morphology of the host galaxies of obscured AGN.

As a final remark we note that, while type 2 quasars are probably numerous, it is likely that they have been elusive so far because have been searched for in the optical rather than in X-rays. In most luminous and/or distant absorbed quasars the Narrow Lines Region may be also obscured or lacking altogether (NGC 6240 is a classical example [22]), and therefore they are missed in optical spectroscopic surveys. Deep **Chandra** and **XMM** surveys will hopefully settle this long-standing issue, and then provide a key test for XRB synthesis models.

Acknowledgements. We thank the **BeppoSAX** SDC, SOC and OCC teams for the successful operation of the satellite and preliminary data reduction and screening, P. Giommi, R. Maiolino, L.A. Antonelli, S. Molendi, M. Mignoli, R. Gilli, G. Risaliti (the “HELLAS boys”) for the fruitful collaboration, M. Salvati, G.C. Perola and G. Zamorani for useful discussions. Partial support from ASI contract ARS-99-75 and MURST grant Cofin98-02-32 is acknowledged.

References

- [1] Akiyama M., et al., 2000, ApJ in press
- [2] Boella G., et al., 1997, A&AS 122, 299
- [3] Cagnoni I., Della Ceca R., Maccacaro T., 1998, ApJ 493, 54
- [4] Comastri A., et al., 1995, A&A 296, 1
- [5] Comastri A., 2000, Astr. Lett & Comm., submitted
- [6] Della Ceca R., et al., 1999, ApJ 524, 674
- [7] Della Ceca R., et al., 2000, astro-ph/9912016
- [8] Fiore F., et al., 1999, MNRAS 306, L55
- [9] Fiore F., Vignali C., Comastri A., et al., 2000, MNRAS submitted
- [10] Gilli R., Risaliti G., Salvati M., 1999, A&A 347, 424
- [11] La Franca F., et al., 2000, in preparation
- [12] Lehmann I., et al., 1999, A&A in press (astro-ph/9911484)
- [13] Maiolino R., et al., 1998, A&A 338, 781
- [14] Maiolino R., et al., 2000, A&A submitted
- [15] Pompilio F., La Franca F., Matt G., 2000, A&A 353, 440
- [16] Risaliti G., Maiolino R., Salvati M., 1999, ApJ 522, 157
- [17] Setti G., Woltjer L., 1989, A&A 224 L21
- [18] Ueda Y., et al., 1999, ApJ 524, L11
- [19] Vecchi A., Molendi S., Guainazzi M., et al., 1999, A&A 349, L73
- [20] Vignali C., Comastri A., Fiore F., 2000, these proceedings
- [21] Vignali C., et al., 2000, MNRAS in press
- [22] Vignati P., et al., 1999, A&A 349, L57